

A Survey on Assessment of Knowledge Attitude and Practice of ADR Reporting Among Health Care Professionals

Dr. P.A. Sreeja*¹, V.J. Dawn², V.R. Irene³, P.V. Soumya³, K. Surabhi Mohan³

1: Professor, Rungta Institute of Pharmaceutical Sciences and Research, Bhilai, Chattisgarh

2: Associate Professor, Department of Pharmacy Practice, Sanjo College of Pharmaceutical Studies, Palakkad, Kerala

3: Assistant Professor, Department of Pharmacy Practice, Sanjo College of Pharmaceutical Studies, Palakkad, Kerala

3: Assistant Professor, Department of Pharmacy Practice, Sanjo College of Pharmaceutical Studies, Palakkad, Kerala

3: Assistant Professor, Department of Pharmacy Practice, Sanjo College of Pharmaceutical Studies, Palakkad, Kerala

ABSTRACT

Adverse drug reactions (ADRs) contribute serious health consequences and underreporting of ADRs by health care professionals (HCPs) is found to be a major problem due to the lack of awareness. The present study was carried out to evaluate the KAP about ADR reporting and pharmacovigilance among HCPs in a rural area in Palakkad District. A questionnaire-based observational survey was conducted among HCPs of a Community Health Centre and a private hospital in Palakkad District. The responses were analyzed and the results were entered in MS-Excel 2010, Statistical analysis was done using Graph pad Prism and chi-square test was done to find the level of significance. Out of 250 questionnaires circulated, 218 health care professionals positively completed and returned of which 23 were from community health centre and 195 from a private hospital. Out of the total participants 104 (48%) were males and 114 (52%) were females. On analyzing the knowledge among health professionals, we found physicians and pharmacist have better knowledge than other HCPs and was found to have statistical relationship ($p < 0.05$). We found a positive response toward attitude and practice and the major problem of under reporting was lack of time and knowledge. Lack of knowledge is found to be a major issue of underreporting and must be rectified by proper awareness campaigns, continuing education programs etc.

KEYWORDS: ADR (Adverse drug reaction), Pharmacovigilance, Continuing education programs

INTRODUCTION

All drugs available in the market are not completely safe and can cause mild to severe life-threatening ADRs.^[1] ADRs are serious health problems that increase morbidity and mortality. It is a common cause for the increase in both the length of hospital stays as well as the cost of treatment.^[2,3]

The World Health Organization (WHO) defines pharmacovigilance (PV) as "the science and activities related to the detection, understanding, and prevention of side effects or other drug-related problems" (interactions, misuse, medication errors, abuse, overdose)^[4]. Adverse drug reaction (ADR) is defined by the WHO as "a response to a drug which is noxious and unintended, and which occurs at doses normally used in man for the prophylaxis, diagnosis, or therapy of disease, or the modification of physiological function"^[5].

All healthcare facilities must promote PV efforts as they are crucial to ensuring patient safety^[6,7]. The primary target group for the effective implementation of any healthcare initiative, including PV, is the HCPs because they make up most of a health care system's resource^[8]. Doctors, pharmacists, nurses, and other paramedics have a huge responsibility to report ADRs and improve the PV systems^[9].

To encourage healthcare professionals and patients for the safe use of medicine, the Ministry of Health and Welfare, Government of India launched the nationwide Pharmacovigilance Programme of India (PvPI) in the year 2010^[10]. Our study aimed to assess the knowledge attitude and practice of ADR reporting and PV among HCPs as we found most of the studies were done among physicians, postgraduate students etc. but assessing the knowledge among other health care professionals like laboratory technicians, dieticians, social workers etc. were also important as they all are an integral part of health care system.

*1 Corresponding Author: sreejavinod318@gmail.com

MATERIALS AND METHODS

A cross-sectional questionnaire-based study design was adapted for a cross-sectional survey by convenience sampling method. The participant's convenient accessibility and proximity was considered for convenient sampling method [11,12]. The survey was conducted for a period of 3 months from September to November 2022. A predesigned validated self-structured questionnaire was used to collect the details about the knowledge, attitude and practice on ADR reporting and Pharmacovigilance among health care professionals at a Community Health Centre and a Private hospital in Palakkad district.

The purpose of the study was explained to the participants and those willing to participate and returned the fully completed questionnaire were included in the study. The questionnaire was circulated among the health care professionals and the responses were collected.

Validation Of Questionnaire:

The questionnaire was designed after reviewing previous research and review articles. Some questions were added on observing the importance of those questions in our study. The prepared questionnaire was circulated among randomly selected professional in the institution and a medical officer. The responses obtained were analyzed and finalized.

The questionnaire consists of total 20 questions, of which 8 knowledge, 8 attitude and 4 practice based questions were asked. Out of the 8 questions, 7 responses are 'yes' or 'no' and 1 was multiple choice. The second part consists of 7 Attitude based questions whose responses were given as 'positive' and 'negative' and a multiple-choice question for analyzing barriers for reporting of ADRs. The third part comprise of 4 Practice based questions and the responses were given as 'yes', 'no' and 'don't know'. The various types of ADR reported by HCPs were also studied.

The data obtained were analyzed and entered in MS Excel 2010. Descriptive statistics like frequency and percentage were used to explain the result. The appropriate statistical analysis was done by Graph Pad Prism software and the level of significance among the participants were calculated by using chi-square test where the p value <0.05 was significant.

RESULT

Out of 250 questionnaires circulated, 218 health care professionals positively completed and returned back of which 23 were from community health centre and 195 from a private hospital. Out of the total participants 104 (48%) were males and 114 (52%) were females. Among them most of the participants were of age between 26 to 45 years with a mean age \pm SD, 35 ± 8.37 . Of the total HCPs, 24 (11%) doctors, 7 (3.21%) house surgeons, 134 (61.46%) nurses, 17 (7.79%) pharmacists, 14 (6.42%) Lab technicians, 9 (4.12%) radiology staffs, 10 (4.58%) social workers and 3 (1.37%) dieticians (**Table 1**)

Table 1: BASIC CHARACTERISTICS OF THE STUDY POPULATION (n= 218)

<i>Characteristics</i>	<i>No: Of Participants (N=218)</i>	<i>Percentage (%)</i>
Gender		
Male	104	47.7
Female	114	52.3
Age group in years		
≤ 25	16	7.3
26-35	108	49.5
36-45	60	27.5
46-55	29	13.3
> 55	5	2.3
Mean age \pm SD	35 ± 8.37	
Study participants		
Doctors	24	11
House surgeons	7	3.2
Nurses	134	61.5
Pharmacist	17	7.8
Lab technicians	14	6.4
Radiology staffs	9	4.1
Social workers	10	4.6
Dietician	3	1.4

a. Knowledge On ADR Reporting and Pharmacovigilance Among Each Professionals:

Our study compared the KAP among each professional. out of the 8 knowledge-based questions, regarding the knowledge on ADR reporting. The result showed that doctors and HS (93.5%) followed by Pharmacists (76.5%) had answered for the question related to safety of medicines available in market as not safe while half of the enrolled nurses (58.2%) and other HCPs (52.8%) found them safe. For the question 2 about safety of medicines other than allopathy, doctors and HS (83.9%) and pharmacist (64.7%) answered not safe but nurses (79.1%) and other HCPs (80.6%) answered that they are safe. 93.5% of Doctors and HS, 82.4% pharmacists, 50.7% nurses and 47.2% of other HCPs respectively have heard about ADRs. On analysis of whether all drugs can cause ADRs and prevention of ADRs, doctors and HS and pharmacist have profound knowledge while nurses and other HCPs have limited. Doctors and HS (80.6% and 74.2%), pharmacist (74.2 and 70.6%) have heard about ADR reporting and pharmacovigilance, respectively. Our study found to have statistical significance (p -value ≤ 0.05) between each group (**Table 2**).

Table-2: ASSESSMENT OF KNOWLEDGE ABOUT ADR, ADR REPORTING AND PHARMACOVIGILANCE (n= 218)

Sl No.	Questions	Category	Yes n(%)	No n(%)	p value
1	Do you think that all medicines available in market are safe	Doctors + HS	2(6.5)	29(93.5)	<0.001
		Nurses	78(58.2)	56(41.8)	
		Pharmacist	4(23.5)	13(76.5)	
		Other HCP	19(52.8)	17(47.2)	
2	Do you think that drugs belonging to other system of medicine (except allopathy) are safe	Doctors + HS	5(16.1)	26(83.9)	<0.001
		Nurses	106(79.1)	28(20.9)	
		Pharmacist	6(35.3)	11(64.7)	
		Other HCPs	29(80.6)	7(19.4)	
3	Have you heard about ADR	Doctors + HS	29(93.5)	2(6.5)	<0.001
		Nurses	68(50.7)	66(49.3)	
		Pharmacist	14(82.4)	3(17.6)	
		Other HCPs	17(47.2)	19(52.8)	
4	Do you think all drugs can cause ADR	Doctors + HS	27(87.1)	4(12.9)	<0.001
		Nurses	57(42.5)	77(57.5)	
		Pharmacist	12(70.6)	5(29.4)	
		Other HCPs	14(38.9)	22(61.1)	
5	Do you think that ADR can be prevented	Doctors + HS	29(93.5)	2(6.5)	<0.001
		Nurses	62(46.3)	72(53.7)	
		Pharmacist	12(70.6)	5(29.4)	
		Other HCPs	15(41.7)	21(58.3)	
6	Have you ever heard of ADR reporting	Doctors + HS	25(80.6)	6(19.4)	0.002
		Nurses	64(47.8)	70(52.2)	
		Pharmacist	12(70.6)	5(29.4)	
		Other HCPs	16(44.4)	20(55.6)	
7	Have you ever heard about Pharmacovigilance?	Doctors + HS	23(74.2)	8(25.8)	0.002
		Nurses	58(43.3)	76(56.7)	
		Pharmacist	12(70.6)	5(29.4)	
		Other HCPs	14(38.9)	22(61.1)	

On analyzing the knowledge on organization responsible for monitoring and reporting ADRs, both Doctors and pharmacist had equal response of answering CDSCO correctly with 29%. Most of the other participants had opted IMA which shows that they were not aware about CDSCO (Fig. 1).




Fig. 1: Knowledge on Organization to Report ADRS

b. Attitude On ADR Reporting and Pharmacovigilance Among Each Professionals:

Out of the 7 attitude-based questions, most of the health care professionals thinks that pharmacist can only report ADR and ADR reporting should be compulsory and voluntary. They also believe that institutions value ≤ 0.05 (Table 3)

TABLE 3 -ATTITUDE TOWARDS REPORTING ADRS AMONG STUDY PARTICIPANTS (n=218)

Sl. No	Attitude Questions	Category	Positive n (%)	Negative n (%)	p-value
1	Do you think that only pharmacist can report ADR?	Doctors + HS	24(77.4)	7(22.6)	<0.001
		Nurses	88(65.7)	46(34.3)	
		Pharmacist	15(88.2)	2(11.8)	
		Other HCPs	11(30.6)	25(69.4)	
2	Do you think ADR reporting should be compulsory	Doctors + HS	26(83.9)	5(16.1)	<0.001
		Nurses	61(45.5)	73(54.5)	
		Pharmacist	12(70.6)	5(29.4)	
		Other HCPs	15(41.7)	21(58.3)	
3	Whether ADR reporting should be voluntary.	Doctors + HS	26(83.9)	5(16.1)	<0.001
		Nurses	46(34.3)	88(65.7)	
		Pharmacist	10(58.8)	7(41.2)	
		Other HCPs	15(41.7)	21(58.3)	
4	Do you believe that workplace environment should encourage reporting an ADR?	Doctors + HS	27(87.1)	4(12.9)	0.004
		Nurses	113(84.3)	21(15.7)	
		Pharmacist	13(76.5)	4(23.5)	
		Other HCPs	21(58.3)	15(41.7)	
5	Are you interested in participating ADR reporting system.	Doctors + HS	24(77.4)	7(22.6)	0.52
		Nurses	111(82.8)	23(17.2)	
		Pharmacist	16(94.1)	1(5.9)	
		Other HCPs	29(80.6)	7(19.4)	
6	Are you interested in learning/ get training about PV.	Doctors + HS	23(74.1)	8(25.8)	0.61
		Nurses	113(84.3)	21(15.7)	
		Pharmacist	14(82.4)	3(17.6)	
		Other HCPs	30(83.3)	6(16.7)	
7	Do you think that every institution should monitor and document ADR?	Doctors + HS	29(93.5)	2(6.5)	0.87
		Nurses	123(91.8)	11(8.2)	
		Pharmacist	15(88.2)	2(11.8)	
		Other HCPs	32(88.9)	4(11.1)	

Our study participants were interested in learning as well as participating in ADR reporting system. All the participants showed a positive attitude towards institutional documentation system of ADRs. However, on statistical evaluation we couldn't find any significant relation among the groups.

Lack of time and knowledge was found to be the main factor of underreporting ADRs. Doctors and HS [n=17 (55%)] found lack of time as most remarkable factor followed by negligence [n=6 (19%)] in not reporting ADRs. Nurses found lack of knowledge/training [n=104 (78%)] as the prime factor followed by lack of time [n=99 (74%)] for not reporting ADRs. Pharmacist found lack of time [n=8 (47%)] and lack of knowledge/training [n=4 (24%)] as the prominent factor for not reporting ADRs respectively. The other HCPs mostly admits lack of knowledge/training [n=26 (72%)] as the barrier for underreporting of ADRs (Fig. 2).




Fig. 2: Barriers for not reporting ADRs.

c. Practice On ADR Reporting and Pharmacovigilance Among Each Professionals:


During the daily practice doctors and HS (90.3%), dispensing pharmacist (70.6%) and nurses (70.9%), other HCPS like radiology staff and laboratory technicians (38.9%) have encountered ADRs in their practice. Among the study subjects, Doctors, and HS (64.5%), nurses (54.5%) pharmacist (29.4%) and other HCPs (8.3%) have filled the ADR reporting form and reported to the institution. Some doctors reported ADRs directly to zonal ADR monitoring centers. Most of the Doctors, Pharmacists and Nurses admit that the institution submits the ADR reporting form to zonal ADR monitoring centers (Table 4).

TABLE 4 - Practice based questions.

S. No.	Practice Questions	Category	Yes n (%)	No n (%)	Don't know n (%)	p-Value
1.	Do you come across any ADR in practice?	Doctors+Hs	28(90.3)	2(6.5)	1(3.2)	0.001
		Nurses	95(70.9)	28(20.9)	11(8.2)	
		Pharmacist	12(70.6)	3(17.4)	2(11.8)	
		Other HCPs	14(38.9)	15(41.7)	7(19.4)	
2.	Have you ever filled any ADR reporting forms?	Doctors+Hs	20(64.5)	9(29.0)	2(6.5)	<0.001
		Nurses	73(54.5)	35(26.1)	26(19.4)	
		Pharmacist	5(29.4)	8(47.1)	4(23.5)	
		Other HCPs	3(8.3)	17(47.2)	16(44.4)	
3.	Whether your institution is practicing ADR reporting to higher centres?	Doctors+Hs	16(51.6)	7(22.6)	8(25.8)	<0.001
		Nurses	64(47.8)	47(35.1)	23(17.2)	
		Pharmacist	4(23.5)	6(35.2)	7(41.2)	
		Other HCPs	3(8.3)	14(38.9)	19(52.7)	
4.	Does your institution provide information or training on ADR monitoring and reporting?	Doctors+Hs	8(25.8)	15(48.3)	8(25.8)	0.626
		Nurses	47(35.1)	56(41.8)	31(23.1)	
		Pharmacist	6(35.3)	5(29.4)	6(35.3)	
		Other HCPs	13(36.1)	11(30.6)	12(33.3)	

The various types of ADRs reported by HCPs were Serious & life threatening, Severe & cause disability, Moderate & cause difficulty, Mild. The reported ADRs by Doctors and HS were mainly Serious & life threatening [n=20 (65%)] and severe & that caused disability [n=19 (61%)]. Nurses and pharmacist reported mainly those ADRs with moderate and cause difficulty [n=68 (51%)] and [n=4 (24%)] respectively followed by mild [n=73 (54%)] and [n=5 (29%)] respectively (Fig. 3).

Fig. 3 - Represents the Types of ADRs Reported By HCPS.



DISCUSSION

The present observational study aimed on the assessment of knowledge, attitude and practice on ADR, ADR reporting and pharmacovigilance. Our study participants include doctors and HS, nurses, pharmacists, radiology staffs, laboratory technicians, social workers, and dieticians. 87.2% responses were obtained from the total distributed questionnaires which was nearly like previous studies conducted by Hussain *et al*^[11]; Alsaleh *et al*^[13]. Most of our participants were females and the mean age \pm SD was 35 ± 8.37 . Most participants were nurses followed by doctors, pharmacist, laboratory technicians, social workers, radiology staff, HS and dieticians.

On analyzing the knowledge about general safety of medicines, ADR reporting and pharmacovigilance, we found that there is lack of knowledge among HCPs other than Doctors and pharmacist which was similar to a study conducted by Jagjit Singh in which they mentioned that physicians had good knowledge when compared to nurses^[14] Even though many of the health care professionals have heard about ADR, most of them doesn't know about the importance of reporting it. A study conducted by Rasqa Adisa have also mentioned the same in their study^[15]. Various studies were conducted for the assessment of knowledge about ADR reporting and pharmacovigilance in and around world stating the importance of monitoring and assessing the same. A study conducted by Ganesan S *et al.*, also mentioned about the inadequate knowledge of health care professionals about pharmacovigilance^[16,17,18].

On assessing the attitude and practice of the health care professionals, majority had a positive response towards participating in ADR reporting system and recommended to have proper training on how to report an encountered ADR. Conducting more training programs in hospitals can improve knowledge about the ADR reporting centres and availability of reporting forms. Major problem found in our study for under reporting an ADR was lack of knowledge/training and lack of time, which was also found to be common in many of the similar studies. A study conducted by Chetna K. Desai reported that lack of knowledge and availability of ADR reporting forms were the major concern of under reporting ADRs^[19]. The practice of documenting the reported ADRs by the institution should also be encouraged which in our study, the participants responded positively. Reporting all types of encountered ADRs by the health care professionals must be practiced and in our study majority of doctors reported serious life-threatening cases more while nurses and pharmacists had a practice of reporting all types of ADRs especially mild to moderate^[20].



LIMITATIONS:

We have assessed the KAP of health care professionals other than physicians, nurses and pharmacist which were rarely done in previous studies. We felt it important to assess the knowledge of other health care professionals like laboratory technicians, radiology staffs, dieticians, and others as they also come across several ADRs of various severity types in their daily practice. Limitation of our study is the smaller number of study participants which can be biased. The value of the study may be added up with a greater number of participants. Although we asked the involvement of more health care professionals, they were not willing to participate due to their busy schedule.

CONCLUSION

Our study concluded, that other than doctors, healthcare professionals have limited knowledge of ADR reporting and Pharmacovigilance. The results proved the need for more frequent awareness campaigns emphasizing the value of timely reporting of experienced ADRs to healthcare professionals by conducting continuous medical education programs or public education. We recommend the posting of a clinical pharmacist with adequate knowledge for the timely monitoring and reporting of ADRs.

ACKNOWLEDGMENT

Sincere thanks to the Principal, Management of Sanjo College of Pharmaceutical Studies (SCOPS)

CONFLICT OF INTEREST: NIL

FUNDING SOURCES: NIL

REFERENCES

1. Almandil NB. Healthcare professionals' awareness and knowledge of adverse drug reactions and pharmacovigilance. *Saudi Med J*. 2016;37(12):1359-1364. doi:10.15537/smj.2016.12.17059
2. Khan Z, Karataş Y, Kiroğlu O. Evaluation of Adverse Drug Reactions in Paediatric Patients: A Retrospective Study in Turkish Hospital [published correction appears in *Front Pharmacol*. 2021 Dec 14;12:813892]. *Front Pharmacol*. 2021;12:786182. Published 2021 Nov 12. doi:10.3389/fphar.2021.786182
3. Hu W, Tao Y, Lu Y, et al. Knowledge, Attitude and Practice of Hospital Pharmacists in Central China Towards Adverse Drug Reaction Reporting: A Multicenter Cross-Sectional Study. *Front Pharmacol*. 2022; 13:823944. Published 2022 Mar 22. doi:10.3389/fphar.2022.823944
4. World Health Organization (WHO). The safety of medicines in public health programmes: pharmacovigilance an essential tool. 2006
5. Coleman JJ, Pontefract SK. Adverse drug reactions. *Clin Med (Lond)*. 2016;16(5):481-485. doi:10.7861/clinmedicine.16-5-481
6. Chan CL, Ang PS, Li SC. A Survey on Pharmacovigilance Activities in ASEAN and Selected Non-ASEAN Countries, and the Use of Quantitative Signal Detection Algorithms. *Drug Saf*. 2017;40(6):517-530. doi:10.1007/s40264-017-0510-9
7. Lihite RJ, Lahkar M. An update on the Pharmacovigilance Programme of India. *Front Pharmacol*. 2015;6:194. Published 2015 Sep 22. doi:10.3389/fphar.2015.00194
8. Singh J, Singh H, Rohilla R, Kumar R, Gautam CS. Lack of Awareness of Pharmacovigilance among Young Health-care Professionals in India: An Issue Requiring Urgent Intervention. *Int J Appl Basic Med Res*. 2018;8(3):158-163. doi:10.4103/ijabmr.IJABMR_423_17



9. Güner MD, Ekmekci PE. Healthcare professionals' pharmacovigilance knowledge and adverse drug reaction reporting behavior and factors determining the reporting rates. *J Drug Assess.* 2019;8(1):13-20. Published 2019 Jan 5. doi:10.1080/21556660.2019.1566137
10. *Pharmacovigilance Programme of India (PvPI)* [Last accessed on 2017 Nov 24]. Available from: <http://www.ipc.gov.in/PvPI/about.html>.
11. Hussain R, Hassali MA, Hashmi F, Akram T. Exploring healthcare professionals' knowledge, attitude, and practices towards pharmacovigilance: a cross-sectional survey. *J Pharm Policy Pract.* 2021;14(1):5. Published 2021 Jan 4. doi:10.1186/s40545-020-00287-3
12. Elfil M, Negida A. Sampling methods in Clinical Research; an Educational Review. *Emerg (Tehran).* 2017;5(1): e52.
13. Alsaleh FM, Alzaid SW, Abahussain EA, Bayoud T, Lemay J. Knowledge, attitude and practices of pharmacovigilance and adverse drug reaction reporting among pharmacists working in secondary and tertiary governmental hospitals in Kuwait. *Saudi Pharm J.* 2017;25(6):830-837. doi:10.1016/j.jsps.2016.12.004
14. Singh J, Singh H, Rohilla R, Kumar R, Gautam CS. Lack of Awareness of Pharmacovigilance among Young Health-care Professionals in India: An Issue Requiring Urgent Intervention. *Int J Appl Basic Med Res.* 2018;8(3):158-163. doi:10.4103/ijabmr.IJABMR_423_17
15. Adisa R, Omitogun TI. Awareness, knowledge, attitude and practice of adverse drug reaction reporting among health workers and patients in selected primary healthcare centres in Ibadan, southwestern Nigeria. *BMC Health Serv Res.* 2019;19(1):926. Published 2019 Dec 3. doi:10.1186/s12913-019-4775-9
16. Ganesan S, Sandhiya S, Reddy KC, Subrahmanyam DK, Adithan C. The Impact of the Educational Intervention on Knowledge, Attitude, and Practice of Pharmacovigilance toward Adverse Drug Reactions Reporting among Health-care Professionals in a Tertiary Care Hospital in South India. *J Nat Sci Biol Med.* 2017;8(2):203-209. doi:10.4103/0976-9668.210014
17. Seid MA, Kasahun AE, Mante BM, Gebremariam SN. Healthcare professionals' knowledge, attitude and practice towards adverse drug reaction (ADR) reporting at the health center level in Ethiopia. *Int J Clin Pharm.* 2018;40(4):895-902. doi:10.1007/s11096-018-0682-0
18. Gupta SK, Nayak RP, Shivarajani R, Vidyarthi SK. A questionnaire study on the knowledge, attitude, and the practice of pharmacovigilance among the healthcare professionals in a teaching hospital in South India. *Perspect Clin Res.* 2015;6(1):45-52. doi:10.4103/2229-3485.148816
19. Desai CK, Iyer G, Panchal J, Shah S, Dikshit RK. An evaluation of knowledge, attitude, and practice of adverse drug reaction reporting among prescribers at a tertiary care hospital. *Perspect Clin Res.* 2011;2(4):129-136. doi:10.4103/2229-3485.86883
20. Behera MR, Tripathy R, Srivastava V, Das MC. Knowledge, attitude, and practice (KAP) of pharmacovigilance among paediatricians of Odisha and factors related to poor reporting of adverse drug reactions. *J Family Med Prim Care.* 2022;11(7):3524-3527. doi:10.4103/jfmprc.jfmprc_2323_21